

# Two-phase increase in the maximum size of life over 3.5 billion years reflects biological innovation and environmental opportunity

Jonathan L. Payne<sup>a,1</sup>, Alison G. Boyer<sup>b</sup>, James H. Brown<sup>b</sup>, Seth Finnegan<sup>a</sup>, Michał Kowalewski<sup>c</sup>, Richard A. Krause, Jr.<sup>d</sup>, S. Kathleen Lyons<sup>e</sup>, Craig R. McClain<sup>f</sup>, Daniel W. McShea<sup>g</sup>, Philip M. Novack-Gottshall<sup>h</sup>, Felisa A. Smith<sup>b</sup>, Jennifer A. Stempien<sup>i</sup>, and Steve C. Wang<sup>j</sup>

<sup>a</sup>Department of Geological and Environmental Sciences, Stanford University, 450 Serra Mall, Building 320, Stanford, CA 94305; <sup>b</sup>Department of Biology, University of New Mexico, Albuquerque, NM 87131; <sup>c</sup>Department of Geosciences, Virginia Polytechnic Institute and State University, Blacksburg, VA 24061; <sup>d</sup>Museum für Naturkunde der Humboldt-Universität zu Berlin, D-10115, Berlin, Germany; <sup>e</sup>Department of Paleobiology, National Museum of Natural History, Smithsonian Institution, Washington, DC 20560; <sup>f</sup>Monterey Bay Aquarium Research Institute, Moss Landing, CA 95039; <sup>g</sup>Department of Biology, Box 90338, Duke University, Durham, NC 27708; <sup>h</sup>Department of Geosciences, University of West Georgia, Carrollton, GA 30118; <sup>i</sup>Department of Geological Sciences, University of Colorado, Boulder, CO 80309; and <sup>j</sup>Department of Mathematics and Statistics, Swarthmore College, 500 College Avenue, Swarthmore, PA 19081

Edited by James W. Valentine, University of California, Berkeley, CA, and approved November 14, 2008 (received for review July 1, 2008)

**The maximum size of organisms has increased enormously since the initial appearance of life >3.5 billion years ago (Gya), but the pattern and timing of this size increase is poorly known. Consequently, controls underlying the size spectrum of the global biota have been difficult to evaluate. Our period-level compilation of the largest known fossil organisms demonstrates that maximum size increased by 16 orders of magnitude since life first appeared in the fossil record. The great majority of the increase is accounted for by 2 discrete steps of approximately equal magnitude: the first in the middle of the Paleoproterozoic Era ( $\approx 1.9$  Gya) and the second during the late Neoproterozoic and early Paleozoic eras (0.6–0.45 Gya). Each size step required a major innovation in organismal complexity—first the eukaryotic cell and later eukaryotic multicellularity. These size steps coincide with, or slightly postdate, increases in the concentration of atmospheric oxygen, suggesting latent evolutionary potential was realized soon after environmental limitations were removed.**

body size | Cambrian | oxygen | Precambrian | trend

Despite widespread scientific and popular fascination with the largest and smallest organisms and numerous studies of body size evolution within individual taxonomic groups (1–9), the first-order pattern of body size evolution through the history of life has not been quantified rigorously. Because size influences (and may be limited by) a broad spectrum of physiological, ecological, and evolutionary processes (10–16), detailed documentation of size trends may shed light on the constraints and innovations that have shaped life's size spectrum over evolutionary time as well as the role of the body size spectrum in structuring global ecosystems. Bonner (17) presented a figure portraying a gradual, monotonic increase in the overall maximum size of living organisms over the past 3.5 billion years. The pattern appears consistent with a simple, continuous underlying process such as diffusion (18), but could also reflect a more complex process. Bonner, for example, proposed that lineages evolve toward larger sizes to exploit unoccupied ecological niches. For decades, Bonner's has been the only attempt to quantify body size evolution over the entire history of life on Earth, but the data he presented were not tied to particular fossil specimens and were plotted without consistent controls on taxonomic scale against a nonlinear timescale. Hence, we have lacked sufficient data on the tempo and mode of maximum size change to evaluate potential first-order biotic and abiotic controls on organism size through the history of life.

Here, we document the evolutionary history of body size on Earth, focusing on the upper limit to size. Use of maximum size allows us to assess constraints on the evolution of large body size

and avoids the more substantial empirical difficulties in determining mean, median, or minimum size for all life or even for many individual taxa. For each era within the Archean Eon (4,000–2,500 Mya) and for each period within the Proterozoic (2,500–542 Mya) and Phanerozoic (542–0 Mya) eons, we obtained the sizes of the largest known fossil prokaryotes, single-celled eukaryotes, metazoans, and vascular plants by reviewing the published literature and contacting taxonomic experts. Sizes were converted to volume to facilitate comparisons across disparate taxonomic groups (see *Data and Methods*). The database is deposited in Data Dryad ([www.datadryad.org](http://www.datadryad.org)) and can be accessed at <http://hdl.handle.net/10255/dryad.222>.

## Results

The maximum body volume of organisms preserved in the fossil record has increased by  $\approx 16$  orders of magnitude over the last 3.5 billion years (Fig. 1). Increase in maximum size occurred episodically, with pronounced jumps of approximately 6 orders of magnitude in the mid-Paleoproterozoic ( $\approx 1.9$  Gya) and during the Ediacaran through Ordovician (600–450 Mya). Thus,  $\approx 75\%$  of the overall increase in maximum body size over geological time took place during 2 geologically brief intervals that together comprise  $<20\%$  of the total duration of life on Earth.

Paleoproterozoic size increase occurs as a single step in Fig. 1, reflecting the presence of *Grypania spiralis* in the Paleoproterozoic (Orosirian) Negaunee Iron Formation of Canada (19). The taxonomic affinities of these fossils are controversial: Their morphological regularity and size suggest they are the remains of eukaryotic organisms (19, 20), but they have also been interpreted as composite microbial filaments (21). Possible trace fossils of similar age and comparable size occur in the Stirling Quartzite of Australia and the Chorhat Sandstone of India, suggesting Orosirian size increase may not have been confined to *Grypania* (22, 23). Slightly younger specimens of similar sizes from the Changzhougou and Changlinggou formations of China

Author contributions: J.L.P., A.G.B., J.H.B., S.F., M.K., R.A.K., S.K.L., C.R.M., D.W.M., P.M.N.-G., F.A.S., J.A.S., and S.C.W. designed research, performed research, analyzed data, and wrote the paper.

The authors declare no conflict of interest.

This article is a PNAS Direct Submission.

Freely available online through the PNAS open access option.

<sup>1</sup>To whom correspondence should be addressed. E-mail: [jlpayne@stanford.edu](mailto:jlpayne@stanford.edu).

This article contains supporting information online at [www.pnas.org/cgi/content/full/0806314106/DCSupplemental](http://www.pnas.org/cgi/content/full/0806314106/DCSupplemental).

© 2008 by The National Academy of Sciences of the USA





volume, associated errors are unlikely to bias results, considering the reported size trends span 16 orders of size magnitude. The observed data and the geometric approximations used to calculate volume for individual fossil specimens are included as Table S1.

Although our database contains only 93 recorded observations, the amount of implicit information recorded is much larger. For example, it is widely agreed that a specimen of *Parapuzosia seppenradensis* is the largest ammonite fossil ever collected (60), and thus, each of the thousands (if not millions) of ammonites ever seen in the field or collected for study must have been smaller than this specimen. By extension of this argument, the database places upper bounds on the sizes of many millions of fossil specimens collected over the past several centuries.

- Alroy J (1998) Cope's rule and the dynamics of body mass evolution in North American fossil mammals. *Science* 280:731–734.
- Hunt G, Roy K (2006) Climate change, body size evolution, and Cope's rule in deep-sea ostracodes. *Proc Natl Acad Sci USA* 103:1347–1352.
- Jablonski D (1997) Body-size evolution in Cretaceous molluscs and the status of Cope's rule. *Nature* 385:250–252.
- Smith FA, et al. (2004) Similarity of mammalian body size across the taxonomic hierarchy and across space and time. *Am Nat* 163:672–691.
- Brown JH, Nicoletto PF (1991) Spatial scaling of species composition—Body masses of North American land mammals. *Am Nat* 138:1478–1512.
- Payne JL (2005) Evolutionary dynamics of gastropod size across the end-Permian extinction and through the Triassic recovery interval. *Paleobiology* 31:269–290.
- Niklas KJ, Enquist BJ (2001) Invariant scaling relationships for interspecific plant biomass production rates and body size. *Proc Natl Acad Sci USA* 98:2922–2927.
- Novack-Gottshall PM, Lanier MA (2008) Scale-dependence of Cope's rule in body size evolution of Paleozoic brachiopods. *Proc Natl Acad Sci USA* 105:5430–5434.
- Huntley JW, Xiao S, Kowalewski M (2006) 1.3 billion years of acritarch history: An empirical morphospace approach. *Precambrian Res* 144:53–68.
- Brown JH (1995) *Macroecology* (Univ of Chicago Press, Chicago).
- Peters RH (1983) *The Ecological Implications of Body Size* (Cambridge Univ Press, New York).
- Calder WA (1984) *Size, Function, and Life History* (Harvard Univ Press, Cambridge, MA).
- Schmidt-Nielsen K (1984) *Scaling, Why is Animal Size so Important?* (Cambridge Univ Press, New York).
- Bonner JT (2006) *Why Size Matters: From Bacteria to Blue Whales* (Princeton Univ Press, Princeton).
- Martin AP, Palumbi SR (1993) Body size, metabolic rate, generation time, and the molecular clock. *Proc Natl Acad Sci USA* 90:4087–4091.
- Jablonski D (1996) in *Evolutionary Paleobiology*, eds Erwin DH, Jablonski D, Lipps JH (Univ of Chicago Press, Chicago), pp 256–289.
- Bonner JT (1988) *The Evolution of Complexity by Means of Natural Selection* (Princeton Univ Press, Princeton).
- McShea DW (1994) Mechanisms of large-scale evolutionary trends. *Evolution (Lawrence, Kans)* 48:1747–1763.
- Han T-M, Runnegar B (1992) Megascopic eukaryotic algae from the 2.1-billion-year-old Negaunee Iron-Formation, Michigan. *Science* 257:232–235.
- Knoll AH, Javaux EJ, Hewitt D, Cohen P (2006) Eukaryotic organisms in Proterozoic oceans. *Philos Trans R Soc London Ser B* 361:1023–1038.
- Samuelsson J, Butterfield NJ (2001) Neoproterozoic fossils from the Franklin Mountains, northwestern Canada: Stratigraphic and palaeobiological implications. *Precambrian Res* 107:235–251.
- Bengtson S, Rasmussen B, Krapež B (2007) The Paleoproterozoic megascopic Stirling biota. *Paleobiology* 33:351–381.
- Seilacher A, Bose PK, Pflüger F (1998) Triploblastic animals more than 1 billion years ago: Trace fossil evidence from India. *Science* 282:80–83.
- Xiao S, Dong L (2006) in *Neoproterozoic Geobiology and Paleobiology*, eds Xiao S, Kaufman AJ (Springer, Dordrecht, The Netherlands), pp 57–90.
- Lamb DM, Awramik SM, Zhu S (2007) Paleoproterozoic compression-like structures from the Changzhongou Formation, China: Eukaryotes or clasts? *Precambrian Res* 154:236–247.
- Kumar S (1995) Megafossils from the Mesoproterozoic Rohtas Formation (the Vindhyan Supergroup), Katni area, central India. *Precambrian Res* 72:171–184.
- Schulz HN, Jørgensen BB (2001) Big bacteria. *Annu Rev Microbiol* 55:105–137.
- Droser ML, Bottjer DJ (1993) Trends and patterns of Phanerozoic ichnofabrics. *Annu Rev Earth Planet Sci* 21:205–225.
- Stanley SM (1973) An explanation for Cope's Rule. *Evolution (Lawrence, Kans)* 27:1–26.
- Gould SJ (1988) Trends as changes in variance—A new slant on progress and directionality in evolution. *J Paleontol* 62:319–329.
- Brocks JJ, Logan GA, Buick R, Summons RE (1999) Archean molecular fossils and the early rise of eukaryotes. *Science* 285:1033–1036.
- Dutkiewicz A, Volk H, George SC, Ridley J, Buick R (2006) Biomarkers from Huronian oil-bearing fluid inclusions: An uncontaminated record of life before the Great Oxidation Event. *Geology* 34:437–440.
- Rasmussen B, Fletcher IR, Brocks JJ, Kilburn MR (2008) Reassessing the first appearance of eukaryotes and cyanobacteria. *Nature* 455:1101–1104.
- Butterfield NJ (2000) *Bangiomorpha pubescens* n. gen., n.sp.: Implications for the evolution of sex, multicellularity, and the Mesoproterozoic/Neoproterozoic radiation of eukaryotes. *Paleobiology* 26:386–404.
- Papineau D, Mojzsis SJ, Schmitt AK (2007) Multiple sulfur isotopes from Paleoproterozoic Huronian interglacial sediments and the rise of atmospheric oxygen. *Earth Planet Sci Lett* 255:188–212.
- Fike DA, Grotzinger JP, Pratt LM, Summons RE (2006) Oxidation of the Ediacaran Ocean. *Nature* 444:744–747.
- Holland HD (2006) The oxygenation of the atmosphere and oceans. *Philos Trans R Soc London Ser B* 361:903–915.
- Canfield DE (2005) The early history of atmospheric oxygen: Homage to Robert M. Garrels. *Annu Rev Earth Planet Sci* 33:1–36.
- Scott C, et al. (2008) Tracing the stepwise oxygenation of the Proterozoic ocean. *Nature* 452:456–459.
- Canfield DE, Poulton SW, Narbonne GM (2007) Late-Neoproterozoic deep-ocean oxygenation and the rise of animal life. *Science* 315:92–95.
- Kaiho K (1998) Global climatic forcing of deep-sea benthic foraminiferal test size during the past 120 m.y. *Geology* 26:491–494.
- Chapelle G, Peck LS (1999) Polar gigantism dictated by oxygen availability. *Nature* 399:114–115.
- McClain CR, Rex M (2001) The relationship between dissolved oxygen concentration and maximum size in deep-sea turrid gastropods: An application of quantile regression. *Mar Biol* 139:681–685.
- Runnegar B (1991) Precambrian oxygen levels estimated from the biochemistry and physiology of early eukaryotes. *Palaogeogr Palaeoclimatol Palaeoecol* 71:97–111.
- Cloud P (1972) A working model of the primitive Earth. *Am J Sci* 272:537–548.
- Runnegar B (1982) The Cambrian explosion: Animals or fossils? *Aust J Earth Sci* 29:395–411.
- Knoll AH, Holland HD (1995) in *Effects of Past Global Change on Life*, ed Commission on Geosciences, Environment and Resources (National Academy Press, Washington, DC), pp 21–33.
- McFadden KA, et al. (2008) Pulsed oxidation and biological evolution in the Ediacaran Doushantuo Formation. *Proc Natl Acad Sci USA* 105:3197–3202.
- Runnegar B (1982) Oxygen requirements, biology and phylogenetic significance of the late Precambrian worm *Dickinsonia*, and the evolution of the burrowing habit. *Alcheringa* 6:223–239.
- Shen Y, Zhang T, Hoffman PF (2008) On the coevolution of Ediacaran oceans and animals. *Proc Natl Acad Sci USA* 105:7376–7381.
- Raff RA, Raff EC (1970) Respiratory mechanisms and the metazoan fossil record. *Nature* 228:1003–1005.
- Acquisti C, Kleffe J, Collins S (2007) Oxygen content of transmembrane proteins over macroevolutionary time scales. *Nature* 445:47–52.
- Fischer WW, Pearson A (2007) Hypotheses for the origin and early evolution of triterpenoid cyclases. *Geobiology* 5:19–34.
- Raymond J, Blankenship RE (2004) Biosynthetic pathways, gene replacement and the antiquity of life. *Geobiology* 2:199–203.
- Chapman DJ, Schopf JW (1983) in *Earth's Earliest Biosphere: Its Origin and Evolution*, ed Schopf JW (Princeton Univ Press, Princeton), pp 302–320.
- Berner RA, Beerling DJ, Dudley R, Robinson JM, Wildman RA (2003) Phanerozoic atmospheric oxygen. *Annu Rev Earth Planet Sci* 31:105–134.
- Geigenberger P (2003) Response of plant metabolism to too little oxygen. *Curr Opin Plant Biol* 6:247–256.
- McShea DW (2001) The hierarchical structure of organisms: A scale and documentation of a trend in the maximum. *Paleobiology* 27:405–423.
- Novack-Gottshall PM (2008) Using simple body size metrics to estimate fossil body volume: Empirical validation using diverse Paleozoic invertebrates. *Palaios* 23:163–173.
- Stevens GR (1988) in *Cephalopods Past and Present*, eds Wiedmann J, Kullmann J (Schweizerbart'sche, Stuttgart, Germany), pp 141–166.